

Application Instructions for Lee Silsby DMSA Transdermal Cream

DMSA Transdermal Cream must be stored at temperatures **BELOW 75° Fahrenheit**. However, do not store in the refrigerator **unless** the room temperature is above 75° Fahrenheit.

1. Remove the blue cap from the tip of the amber syringe containing the DMSA cream.
2. Squirt the instructed amount of cream onto thin-skinned areas of the patient's body such as the tops of feet, thighs (from the knee directly up, not the inner thigh), ankles, shins, inner forearm or lower back. **Do not yet rub in until the steps below are completed.**
3. Next, remove the small cap from the top of the EMP jar labeled **IPM Cream for DMSA**.
4. Using your thumb, push up from the bottom of the jar until an amount approximately equal to the DMSA cream is squeezed out of the top of the jar. Apply on top of, or directly next to, the DMSA cream.
5. Wearing a glove or finger cot, rub and mix **both** creams together. Latex gloves are fine to use as long as your child does not have a sensitivity to latex. If such a sensitivity does exist, please use Nitrile gloves, which are latex-free.
6. If possible, rotate the site of application each time the DMSA is applied.
7. If you will be ordering **another cycle of DMSA**, please call just after beginning your second round of application so that the new prescription will reach you in a timely fashion.

Please check with your practitioner about adequate mineral supplementation during detoxification, or call us with any questions, at: 216-321-4300 or 800-918-8831

Alan Israel, R.Ph.
Robert Wright, R.Ph.

Visit our website for new updates and information at www.leesislby.com

Not all DMSA is made the same: Unless properly formulated, DMSA degrades rapidly and loses its potency. Our formula and two-part application system enhances the stability of DMSA, keeping odor to a minimum, and ensuring that the last dose is as effective as the first.


Please see reverse side for instructions on how to use your EMP jar.

Using Your EMP Jar


The EMP jar is an effective and easy way to apply your medicine. Just follow these three simple steps.



1. Remove the **white** cap from the top of the EMP jar. **DO NOT** unscrew the lid, which will be red, blue or white in color.



2. Hold the EMP jar upright. Using your thumb, push **lightly** on the white circle located in the center of the bottom of the EMP jar until the desired amount of medication coming out of the top of EMP jar is visible.



3. Using your finger, wipe the medication from the top of the EMP jar and apply it to the specified area. If the medication is being applied by someone other than patient, a finger cot or latex gloves are recommended, so long as the patient does not have a sensitivity to latex. If such a sensitivity does exist, please use Nitrile gloves, which are latex-free.